

Probing the Enhancon via Calculations in Supersymmetric Gauge Theory

Steven Corley and David A. Lowe

Department of Physics
Brown University
Providence, RI 02912
scorley, lowe@het.brown.edu

Abstract

We consider the $\mathcal{N} = 2$ gauge theory on N D7-branes wrapping $K3$, with D3-brane probes. In the large N limit, the D7-branes blow up to form an enhancon shell. We probe the region inside and outside the enhancon shell using the D3-branes, and compute the probe metric using the Seiberg-Witten formalism. Supergravity arguments suggest a flat interior up to $1/N$ corrections, and indeed our results for the D3-brane probes are consistent with that. By including the dynamics of the branes, these results, together with those of hep-th/0204050, demonstrate the robustness of the enhancon mechanism beyond patching together of supergravity solutions with D-brane source junction conditions.

1 Introduction

One of the most important questions that a consistent theory of quantum gravity should answer is how spacetime singularities are to be resolved. This is of particular importance in cosmology, where we believe the big bang emerged from a spacelike or null singularity. In the present paper we will consider the resolution of time-like singularities in string theory using the so-called enhancon mechanism. Such singularities are presently under far better calculation control than the singularities of interest in cosmology, because one may use techniques from supersymmetric gauge theory.

The enhancon mechanism is the way string theory resolves a particular type of timelike singularity associated with a stack of D-branes wrapping a compact cycle of the internal compactified space [1]. The would-be timelike singularity of the naive supergravity solution is resolved by brane sources expanding out to form a shell. Inside the shell of brane sources, it has been argued the spacetime is flat.

The main motivation for the enhancon picture comes from studying the behavior of test brane probes. In [1], a configuration of coincident D-branes wrapping a $K3$ surface was considered, and this was probed by a test D-brane also wrapping the $K3$. The coefficient of the v^2 term in the effective action for the probe changes sign at a point of order a string length away from a would-be naked singularity in the naive supergravity solution, where the volume of the $K3$ reduces to the self-dual point. This coefficient of the v^2 term can be thought of as inertial tension. The change in sign of the inertial tension suggests the following self-consistent picture: the wrapped D-branes expand out to form an enhancon shell, and incoming probe D-branes spread out as they approach the shell and dissolve into it as their inertial tension vanishes. In particular, the region inside the enhancon shell cannot be probed by wrapped D-branes. In earlier work [2], we computed the moduli space metric for such wrapped probe branes for the case of D7-branes, by performing calculations using the low energy effective action of the $\mathcal{N} = 2$ supersymmetric $SU(N)$ gauge theory. This low energy description of the dynamics is appropriate when the probe branes are within a string length of the enhancon shell. At large N , we found agreement with supergravity predictions, but we also obtained $1/N$ corrections, as well as instanton corrections nonperturbative in $1/N$. See [3, 4] for other studies of the enhancon from the gauge theory perspective.

An unwrapped D-brane probe has a rather different experience as it approaches the enhancon shell. For the remainder of this paper we will restrict our attention to an enhancon configuration built out of D7-branes wrapping a $K3$, and will add unwrapped D3-branes as probes. Since the D3-brane does not wrap the $K3$, its inertial tension remains non-zero as it passes through the enhancon shell [5]. The main focus of the present paper will be to use such probes to examine the interior of the enhancon shell. We will solve for the low-energy effective action of such probes using techniques from $\mathcal{N} = 2$ supersymmetric gauge theory [6, 7].

These interior probes are interesting, because the region inside need not be governed by low energy supergravity. Of course if one assumes the low energy supergravity equations are valid in the interior, flat space is guaranteed by Gauss' law. However this assumption

need not be correct a priori. From the point of view of solving the low energy equations of motion, with boundary conditions placed at spatial infinity, the interior of the enhancon is inside a shell where stringy physics is important since the curvature becomes of order the string scale near the shell. So the possibility that stringy physics is important in the interior cannot be ruled out without further consideration. We will find that curvature in the interior is actually of order N^{-2} due to back-reaction effects.

Furthermore the patching of supergravity solutions considered in [1, 5] relies heavily on the presence of unbroken supersymmetry, so it is conceivable the solution is unstable to generic perturbations. The enhancon shell is the minimal radius at which D-brane source boundary conditions can be consistently placed, but any larger radius will also do, so the configuration is at best marginally stable [5]. By exactly solving the low-energy effective action for probe branes in this background, we are able to take into account effects of interactions and small deviations away from supersymmetry. Our results place the enhancon mechanism on a much more robust footing.

At the same time, the gauge theory results also highlight the limitations of the enhancon mechanism. Once one gives up spherical symmetry in the large N limit, one can try to consider spacetime singularities dual to Argyres-Douglas [8] points in the moduli space of the gauge theory. At these points an enhancon mechanism does not appear to work [2]. It remains an interesting open problem to understand the spacetime physics associated with these singularities.

A summary of the layout of the paper is as follows. In section 2 we review the supergravity solution for D7-branes wrapping $K3$ and the enhancon mechanism in this context. In section 3 we probe the enhancon using a pair of unwrapped D3-branes by solving for the low energy effective action of $\mathcal{N} = 2$ supersymmetric $SU(N)$ gauge theory with two fundamental hypermultiplets. In section 4 we briefly consider the case of a single hypermultiplet, which is technically more complicated. In section 5, we use these results to compute the moduli space metric for the probe, and compare to supergravity expectations. Section 6 contains some conclusions and prospects for the future.

2 Supergravity solution

We consider the supergravity solution for N D7-branes wrapping $K3$ [1]. The spacetime solution in string metric is

$$\begin{aligned} ds^2 &= Z_3^{-1/2} Z_7^{-1/2} \eta_{\mu\nu} dx^\mu dx^\nu + Z_3^{1/2} Z_7^{1/2} (\alpha')^2 du d\bar{u} + (2\pi R)^2 Z_3^{1/2} Z_7^{-1/2} ds_{K3}^2 \\ e^\Phi &= g Z_7^{-1} \\ Z_3 &= \frac{gN}{2\pi} \frac{(\alpha')^2}{R^4} \ln(U/\rho_3) \\ Z_7 &= \frac{gN}{2\pi} \ln(\rho_7/U) , \end{aligned} \tag{1}$$

where $U = |u|$, $V = (2\pi R)^4$ is the $K3$ volume and g is the string coupling. Here $\eta_{\mu\nu}$ is the Minkowski metric, and ds_{K3}^2 is the $K3$ metric. The constants ρ_3 and ρ_7 are new

constants of integration that appear in the supergravity solution. For $N \leq 24$ this type of configuration can be realized in F-theory. However for $N > 24$ this supergravity solution must be embedded in some yet to be discovered generalization of F-theory. This solution has a singularity at $U = \rho_7$ where the dilaton blows up and the supergravity description breaks down. $U = \rho_3$ is the repulson singularity (we assume $\rho_3 < U < \rho_7$), where the dilaton remains finite, but the curvature blows up. This singularity is cutoff by the enhancon mechanism, with the enhancon shell sitting just outside $U = \rho_3$.

For a wrapped D7-brane probe, the inertial tension is

$$\frac{1}{2g}(\mu_7 V Z_3 - \mu_3 Z_7) , \quad (2)$$

where $\mu_3 = (2\pi)^{-3}\alpha'^{-2}$ and $\mu_7 = (2\pi)^{-7}\alpha'^{-4}$. This changes sign as discussed above, as U approaches ρ_3 . This is the sign of the instability leading to the expansion of the brane charge into the enhancon shell. The inertial tension vanishes at the enhancon radius. Because the Compton wavelength of these branes becomes large near the enhancon shell, we cannot use these branes as probes of the interior of the enhancon.

For an unwrapped D3-brane, the inertial tension is

$$\frac{1}{2g}\mu_3 Z_7 , \quad (3)$$

As remarked above, this is finite at the enhancon radius $U = U_e$, and remains finite inside the enhancon shell, where Z_7 is to be replaced by $Z_7(U_e)$. These branes can be used as probes of the interior of the enhancon shell, where they simply see a flat moduli space metric.

3 Probing the enhancon

We begin by considering the case of two D3-brane probes in the limit that they approach the collection of D7-branes. As we will see, considering two probe branes leads to technical simplifications that allow us to obtain explicit results for the period matrix. We will work in a limit where the inertial tension of the D3-branes is much smaller than that of the N wrapped D7-branes so that they may be considered true test probes of the enhancon geometry. In this limit the dynamics reduces to simply $SU(N)$ with two fundamental hypermultiplets, rather than the full $SU(N) \times U(2)$ gauge theory with a bifundamental $(N, 2)$ hypermultiplet. We will comment further on this decoupling later. As is now well-known, the low energy effective action can be obtained from the following Riemann curve [9, 10, 11]

$$Y^2 = A(X)^2 - B(X) \quad (4)$$

with $A(X) = \prod_{i=1}^N (X - \phi_i)$ and $B(X) = \Lambda^{2N-n_3} \prod_{f=1}^{n_3} (X + m_f)$ where the QCD scale is Λ and for us $n_3 = 2$ with m_f the masses of each hypermultiplet, and the ϕ_i are the classical moduli space coordinates corresponding to the transverse positions of the D7-branes.

For simplicity, we take all of the classical moduli space coordinates $\phi_i = 0$ as this is the point that corresponds to the spherically symmetric enhancon [1, 2]. To investigate the physics inside the enhancon we take the masses of the hypermultiplets to be small, i.e., $m_1/\Lambda, m_2/\Lambda \ll 1$. Naively this is the correct limit to take as these masses measure the length of the strings stretched between the constituent D7-branes and the probe D3-branes. The branch points separate into two classes. Most $(2N - 2)$ of the branch points lie on a circle (the enhancon) about the origin. In a double expansion in m_1/Λ and m_2/Λ these branch points are given by

$$\begin{aligned} X_k &= X_k^{(0)}(1 + X_k^{(1)} + X_k^{(2)} + \dots) \\ X_k^{(0)} &= \Lambda e^{i\pi(k-1)/(N-1)} \\ X_k^{(1)} &= \frac{\Lambda^{2N-2}(m_1 + m_2)}{2N(X_k^{(0)})^{2N-1} - 2\Lambda^{2N-2} X_k^{(0)}} \\ X_k^{(2)} &= \frac{\Lambda^{2N-2}((X_k^{(0)} X_k^{(1)})^2 + (m_1 + m_2)X_k^{(0)} X_k^{(1)} + m_1 m_2) - N(2N - 1)(X_k^{(0)})^{2N}(X_k^{(1)})^2}{2N(X_k^{(0)})^{2N} - 2\Lambda^{2N-2}(X_k^{(0)})^2} \end{aligned} \quad (5)$$

for $1 \leq k \leq 2(N - 1)$. The remaining two roots lie near the center of the circle and are given approximately by

$$\begin{aligned} X_{m_1} &= -m_1 \left(1 - \frac{1}{2} \left(1 - \frac{m_2}{m_1} \right) - \frac{1}{2} \sqrt{\left(1 - \frac{m_2}{m_1} \right)^2 + 4 \left(\frac{m_1}{\Lambda} \right)^{2(N-1)} + \dots} \right) \\ X_{m_2} &= -m_2 \left(1 - \frac{1}{2} \left(1 - \frac{m_1}{m_2} \right) + \frac{1}{2} \sqrt{\left(1 - \frac{m_1}{m_2} \right)^2 + 4 \left(\frac{m_2}{\Lambda} \right)^{2(N-1)} + \dots} \right) \end{aligned} \quad (6)$$

We have assumed that $m_1 \leq m_2$ in writing the above form of the roots. Moreover when $m_1 = m_2$ we shall always take the root X_{m_1} to satisfy $|X_{m_1}| < |X_{m_2}|$ (which is the case if the limit is taken on the expressions given).

To evaluate the periods we take the following basis of cycles. The α_k cycles encircle the X_{2k} and X_{2k+1} branch points for $1 \leq k \leq (N - 2)$ and the α_{N-1} cycle encircles X_1 and X_{m_1} , the γ_j cycles encircle the X_{2j-1} and X_{2j} branch points for $1 \leq j \leq (N - 1)$, and the β_j cycles are defined by

$$\begin{aligned} \beta_j &\equiv \gamma_N + \sum_{k=1}^j \gamma_k = - \sum_{k=j+1}^{N-1} \gamma_k, \quad 1 \leq j \leq (N - 2) \\ &\equiv \gamma_N = - \sum_{k=1}^{N-1} \gamma_k, \quad j = (N - 1). \end{aligned} \quad (7)$$

One needs to be careful in deforming contours as we have in the above expressions. We will comment shortly on why this is valid. One can easily check that the α and β cycles form a canonical basis.

Before continuing it is useful to discuss in a little more detail figure 1. Roughly speaking a pair of branch points corresponds to a D7-brane. One has to be a little careful




Figure 1: Diagram of the branch points and cycles for the two hypermultiplet case.

with such statements, especially in the context here where we are considering a large number of branch points ($2N$ with large N) that are closely packed together, and moreover because a given pair of branch points could actually correspond to more than just a single D7-brane. At any rate, given this identification, figure 1 shows that there is a single D7-brane lying near the center of the enhancon. What has happened is the following. Consider first the case where the hypermultiplet masses are large, $m_1/\Lambda, m_2/\Lambda \gg 1$. In this limit the branch points of the Seiberg-Witten curve are given approximately by $X_k = (\Lambda^{2(N-1)}m_1m_2)^{1/2N}e^{i\pi k/N}$ for $1 \leq k \leq 2N$, i.e., all branch points are on the enhancon circle so that all D7-branes are on the enhancon circle. In this limit one could integrate out the hypermultiplets to arrive at the pure gauge theory at the enhancon point of the moduli space. As the masses of the hypermultiplets are decreased, the size of the enhancon shrinks and moreover a pair of branch points moves toward the center of the circle. In the limit that both masses tend to zero, these two branch points merge at the origin and the low-energy effective action breaks down. What we will now show is that in the limit where both masses are taken to zero, certain charged states associated to fundamental hypermultiplets, and corresponding to the β_{N-1} (or γ_N) contour, are becoming massless. In other words, the D7-brane that lies inside the enhancon is really a bound state of the D7 and D3-branes.

The periods are given by integrating the Seiberg-Witten form $d\lambda$ around the basis of contours described above and depicted in figure 1. Specifically one has

$$\begin{aligned} a_j &= \frac{1}{2\pi i} \oint_{\beta_j} d\lambda \\ a_{D_k} &= \frac{1}{2\pi i} \oint_{\alpha_k} d\lambda \end{aligned} \tag{8}$$

where $d\lambda$ is given by [11, 10, 9]

$$d\lambda = \frac{XdX}{Y} \left(\mathcal{C}'(X) - \mathcal{C}(X) \frac{G'(X)}{2G(X)} \right), \quad (9)$$

and $\mathcal{C}(X) = \prod_{i=1}^N (X - \phi_i)$ and $G(X) = \prod_{l=1}^{N_f} (X + m_l)$. In our case $\mathcal{C}(X) = X^N$ and $G(X) = (X + m_1)(X + m_2)$. It is important to note that the $G'(X)/G(X)$ term in $d\lambda$ has poles at $X = -m_1$ and $X = -m_2$. We take our contours above so as to not encircle these poles. The poles nevertheless will contribute to the mass formula as will be seen below.

To evaluate the periods on the enhancon it is convenient to parametrize the integrals in terms of a new variable φ as

$$X(\varphi) = X_k|_{k \rightarrow k + \varphi/\pi}. \quad (10)$$

The usefulness of this parametrization lies in the fact that Y is given exactly by

$$Y(\varphi) = \Lambda^N e^{i\pi k/(N-1)} e^{i\varphi/(N-1)} \sqrt{e^{i2\varphi} - 1}. \quad (11)$$

As a consequence, substituting the expansion (10) into the Seiberg-Witten form (9) one finds that all φ integrals reduce to the form

$$\begin{aligned} F(n) &= \int_0^\pi d\varphi \frac{e^{i(n-1)\varphi/(N-1)}}{\sqrt{e^{i2\varphi} - 1}} = \frac{e^{-i\pi/4}}{\sqrt{2}} \int_0^\pi d\varphi \frac{e^{i((n-1)/(N-1)-1/2)\varphi}}{\sqrt{\sin(\varphi)}} \\ &= -i2\sqrt{\pi} \frac{N-1}{n-1} \frac{\Gamma(1 + (n-1)/(2N-2))}{\Gamma(1/2 + (n-1)/(2N-2))} e^{in\pi/(2N-2)} \sin\left(\frac{(n-1)\pi}{2N-2}\right). \end{aligned} \quad (12)$$

This leads to the expressions for the $\tilde{a}_j$ (corresponding to the γ_j contours) and a_{D_k} periods

$$\begin{aligned} \tilde{a}_j &= \frac{\Lambda}{\pi(1-1/N)} \left(e^{i2\pi N(j-1)/(N-1)} F(N+1) \left(1 - \frac{1}{N} \right) \right. \\ &\quad \left. - \frac{m_1 + m_2}{2\Lambda} e^{i2\pi(j-1)} \left(\frac{F(-N+2)}{N-1} - \frac{F(N)}{N} \right) + \dots \right), \quad 1 \leq j \leq (N-1) \\ a_{D_k} &= \tilde{a}_j|_{j \rightarrow k+1/2}, \quad 1 \leq k \leq (N-2). \end{aligned} \quad (13)$$

To higher order we show in the appendix that the expansion of the periods in terms of the mass parameter m defined as $m \equiv m_1$ with $\xi \equiv m_2/m_1$ is given by

$$\tilde{a}_j = \sum_{n=0}^{\infty} c_n(\xi) \left(\frac{m}{\Lambda} \right)^n e^{i2\pi(j-1)(1-n)/(N-1)}, \quad (14)$$

with a similar expansion for a_{D_k} .

As an application of this expansion we note that due to the phase factor one obtains the formula for the sum of periods

$$\sum_{j=1}^{N-1} \tilde{a}_j = (N-1) \sum_{p=0}^{\infty} c_{p(N-1)+1}(\xi) \left(\frac{m}{\Lambda} \right)^{p(N-1)+1}, \quad (15)$$

where from (13) we find that $c_1(\xi)$ is given by

$$\begin{aligned} c_1(\xi) &= -\frac{1+\xi}{2\pi(1-1/N)} \left(\frac{F(-N+2)}{(N-1)} - \frac{F(N)}{N} \right) \\ &= \frac{1+\xi}{2(N-1)} , \end{aligned} \quad (16)$$

where we have used $F(-N+2) = 0$ and $F(N) = \pi$ as follows from (13). However the sum (15) corresponds to integrating $d\lambda$ around the contour $\sum_{j=1}^{N-1} \gamma_j$, which can be deformed to the contour $-\gamma_N$ (which by definition does not include the poles at $X = -m_1$ and $X = -m_2$) plus the contribution of the poles of $d\lambda$ at $X = -m_1$ and $-m_2$ plus the contribution due to the pole at infinity. One can show easily however that the contribution from the pole at infinity exactly cancels the contributions of the poles at $X = -m_1$ and $X = -m_2$. So in the end we find that

$$a_{N-1} = \tilde{a}_N = - \sum_{j=1}^{N-1} \tilde{a}_j. \quad (17)$$

Consequently the period a_{N-1} vanishes linearly with m .

Recall that the BPS mass formula [11, 10, 9] is given by

$$M^2 = 2|Z|^2 , \quad (18)$$

where the central charge Z is

$$Z = \sum_{i=1}^N (n_e^i a_i + n_m^i a_{D_i}) + \sum_{j=1}^2 S_j m_j. \quad (19)$$

The constants n_e^i and n_m^i are the electric and magnetic charges respectively and satisfy the constraints $\sum_{i=1}^N n_e^i = 0 = \sum_{i=1}^N n_m^i$. The S_i are the $U(1)$ charges corresponding to global symmetries that are broken by non-zero values of the masses. Also the extra cycles a_N and a_{D_N} are defined by the conditions $\sum_{i=1}^N a_i = 0$ and $\sum_{i=1}^N a_{D_i} = 0$ respectively. The massive vector bosons then carry electric charges $(0, \dots, 0, 1, 0, \dots, 0, -1, 0, \dots, 0)$ while the fundamental matter, quarks, carry electric charges $(0, \dots, 0, 1, 0, \dots, 0) - (1/N, \dots, 1/N)$. It follows from the expression for the a_{N-1} period (17) that its associated quark state is becoming massless in the $m_1, m_2 \rightarrow 0$ limit. Therefore a hypermultiplet is becoming massless, or in the dual string theory description, the probe D7-brane inside the enhancon is really a bound state of the D7 and the D3-branes. To learn more about this D7-D3 bound state, and more about the geometry inside the enhancon, we now compute the period matrix.

To compute the period matrix we must evaluate the periods

$$\frac{\partial a_j}{\partial \xi_n} = \oint_{\beta_j} \omega_n, \quad \frac{\partial a_{Dm}}{\partial \xi_n} = \oint_{\alpha_m} \omega_n , \quad (20)$$

where we define the basis of holomorphic 1-forms by

$$\omega_n \equiv -\frac{1}{2\pi i} \frac{X^{n-1} dX}{Y}, \quad 1 \leq n \leq (N-1). \quad (21)$$

To evaluate the contour integrals along the enhancon circle we again use the parametrization given in (10). To leading order the periods are then given by

$$\begin{aligned} \frac{\partial \tilde{a}_j}{\partial \tilde{\xi}_n} &= -\Lambda^{n-N} \frac{F(n)}{\pi(N-1)} e^{i2\pi(j-1)(n-1)/(N-1)} + \dots, \quad 1 \leq j \leq (N-1) \\ \frac{\partial a_{D_k}}{\partial \tilde{\xi}_n} &= \frac{\partial \tilde{a}_j}{\partial \tilde{\xi}_n} \Big|_{j \rightarrow (k+1/2)}, \quad 1 \leq k \leq (N-2). \end{aligned} \quad (22)$$

The remaining periods are given by the integrals

$$\frac{\partial a_{D_{N-1}}}{\partial \tilde{\xi}_n} = -\frac{i}{\pi} \int_{X_1}^{X_{m_1}} \frac{X^{n-1} dX}{\sqrt{X^{2N} - \Lambda^{2N-2}(X+m_1)(X+m_2)}}. \quad (23)$$

As above we may expand in powers of m_1 and m_2 to obtain to leading order the integral

$$\frac{\partial a_{D_{N-1}}}{\partial \tilde{\xi}_n} = -\frac{i}{\pi} \int_{\Lambda}^0 \frac{X^{n-1} dX}{\sqrt{X^{2N} - \Lambda^{2N-2} X^2}} + \mathcal{O}(m/\Lambda). \quad (24)$$

Redefining the variable of integration by $X = \Lambda e^{x/(N-1)}$, these periods reduce to

$$\begin{aligned} \frac{\partial a_{D_{N-1}}}{\partial \tilde{\xi}_n} &= \frac{i}{\pi(N-1)} \Lambda^{n-N} \int_{-\infty}^0 dx \frac{e^{x(n-(N+1)/2)/(N-1)}}{\sqrt{2 \sinh(x)}} + \mathcal{O}(m/\Lambda) \\ &= \frac{\Lambda^{n-N}}{\sqrt{\pi}} \frac{1}{n-1} \frac{\Gamma(1+(n-1)/(2N-2))}{\Gamma(1/2+(n-1)/(2N-2))} + \mathcal{O}(m/\Lambda), \end{aligned} \quad (25)$$

valid for $n \geq 2$.

The power series expansion in m_1 and m_2 is not valid for the $n=1$ case. In this case one notes that the factor of $\tilde{X}^{2N}$ in the denominator (defining the dimensionless variable $\tilde{X} \equiv X/\Lambda$) can be dropped to leading order since the limits of integration (to leading order) are 0 and 1. In fact this same approximation can be used more generally for $n \ll N$ to obtain the same expression given above. The resulting integral is straightforward to evaluate and we find

$$\frac{\partial a_{D_{N-1}}}{\partial \tilde{\xi}_1} \approx -\frac{\Lambda^{1-N}}{\pi} \ln \left(\frac{m_1}{4\Lambda} \zeta - \frac{1}{2} \sqrt{\left(\frac{m_1 - m_2}{2\Lambda} + \frac{m_1}{2\Lambda} \zeta \right) \left(\frac{m_2 - m_1}{2\Lambda} + \frac{m_1}{2\Lambda} \zeta \right)} \right), \quad (26)$$

where we have defined

$$\zeta \equiv \sqrt{\left(1 - \frac{m_2}{m_1} \right)^2 + 4 \left(\frac{m_2}{\Lambda} \right)^{2(N-1)}}. \quad (27)$$

Some special limiting values for $\partial a_{D_{N-1}}/\partial \xi_1$ are given by

$$\begin{aligned}\frac{\partial a_{D_{N-1}}}{\partial \tilde{\xi}_1} &\approx -\frac{\Lambda^{1-N}}{\pi} \ln\left(\frac{m_2 - m_1}{4\Lambda}\right), \quad \left(\frac{m_1}{\Lambda}\right)^{2(N-1)} \ll \left(1 - \frac{m_2}{m_1}\right)^2 \\ &\approx -\frac{\Lambda^{1-N}}{\pi} N \ln\left(\frac{m_1}{\Lambda}\right), \quad m_1 = m_2.\end{aligned}\tag{28}$$

The period matrix, which is expressed in terms of the periods by

$$\tau_{mj} = \frac{\partial a_{D_m}}{\partial a_j} = \sum_{n=1}^{N-1} \frac{\partial a_{D_m}}{\partial \xi_n} \left(\frac{\partial a}{\partial \xi}\right)_{nj}^{-1},\tag{29}$$

is now straightforward to compute and is

$$\begin{aligned}\tau_{mj} &= \frac{i}{(N-1)} \left(\cot\left(\pi \frac{m-j+1/2}{N-1}\right) - \cot\left(\pi \frac{m-j-1/2}{N-1}\right) \right), \quad 1 \leq m, j \leq (N-2) \\ &= \frac{-i}{(N-1)} \frac{e^{-i\pi(m-1/2)/(N-1)}}{\sin(\pi(m-1/2)/(N-1))}, \quad 1 \leq m \leq (N-2), \quad j = (N-1) \\ &= i\Lambda^{N-1} \frac{\partial a_{D_{N-1}}}{\partial \tilde{\xi}_1}, \quad m = j = (N-1),\end{aligned}\tag{30}$$

where the remaining elements are fixed by symmetry, $\tau_{mj} = \tau_{jm}$.

This is our final result for the period matrix and one of the main results of this paper. Let us now see how to interpret this result. Near the singularity $m_1 = m_2 = 0$ we can use (17) to express $\tau_{N-1, N-1}$ in terms of a_{N-1}

$$\tau_{N-1, N-1} \sim -\frac{i}{\pi} \ln(a_{N-1} + m)\tag{31}$$

in the case $m_1 = m_2 = 2$. This is exactly the singularity we expect when the low energy dynamics reduces to supersymmetric QED with two massless hypermultiplets. We interpret this as a bound state of a D7-brane with the two D3-branes that is able to probe inside the enhancon shell. We will use the results of this section to compute the probe metric in section 5.

4 The single hypermultiplet case

We consider in this section the technically more complicated case of the addition of a single hypermultiplet. The Seiberg-Witten curve becomes

$$Y^2 = X^{2N} - \Lambda^{2N-1}(X + m)\tag{32}$$

where as before we have taken the vevs of the adjoint scalar to be zero, i.e., $\phi_j = 0$. The parameter m is the mass of the hypermultiplet and moreover is the free parameter that we

will vary. In the D-brane picture it corresponds to the mass of a string stretched between the D3-brane probe and the D7-branes.

We consider the case when the D3-brane probe is close to the background D7-branes, i.e., when $m/\Lambda \ll 1$. In this limit the roots to the Seiberg-Witten curve are straightforward to compute in an expansion in powers of m/Λ . In particular one class of roots comes from taking $m = 0$ to leading order to obtain the equation $X^{2N-1} = \Lambda^{2N-1}$. These roots are given by (including some higher order corrections)

$$\begin{aligned} X_k &= X_k^{(0)}(1 + X_k^{(1)} + X_k^{(2)} + \dots) \\ X_k^{(0)} &= \Lambda e^{i2\pi k/(2N-1)} \\ X_k^{(1)} &= \frac{m}{\Lambda} \frac{\Lambda^{2N}}{X_k^{(0)}(2N(X_k^{(0)})^{2N-1} - \Lambda^{2N-1})} \\ X_k^{(2)} &= -N(2N-1) \frac{(X_k^{(1)})^2 (X_k^{(0)})^{2N-1}}{2N(X_k^{(0)})^{2N-1} - \Lambda^{2N-1}} \end{aligned} \quad (33)$$

where $1 \leq k \leq (2N-1)$. The remaining root comes from assuming that $X = -m$ to leading order. Including some correction terms we find

$$X_{2N} = -m \left(1 - \left(\frac{m}{\Lambda} \right)^{2N-1} + \dots \right). \quad (34)$$

To compute the periods a_{D_k} and a_j we define the associated cycles α_k and β_j as follows. The α_k cycle is defined to be the contour encircling the X_{2k+1} and X_{2k+2} branch points for $1 \leq k \leq (N-1)$. The β_j cycles are defined in terms of the γ_k cycles via

$$\beta_j \equiv \sum_{k=1}^j \gamma_k \quad (35)$$

with the γ_k cycle given by the contour that encircles the X_{2k} and X_{2k+1} branch points for $1 \leq k \leq (N-1)$. Figure (2) illustrates the “pac-man” nature of the the choice of branch cuts and cycles described above. We shall denote the periods corresponding to the γ_k cycles by $\tilde{a}_k$.

As in the previous case, we evaluate all periods except for $\tilde{a}_{N-1}$ by parameterizing the X integration variable as

$$X(\varphi) = X_k|_{k \rightarrow k+\varphi/\pi} \quad (36)$$

where the limits of integration for φ are now just 0 and π and k would be $2m+1$ for the α_m cycle and $2j$ for the γ_j cycle. The main advantage to this parametrization is that Y reduces exactly to

$$Y = \Lambda^N e^{i\pi k/(2N-1)} e^{i\varphi/(2N-1)} \sqrt{(e^{i2\varphi} - 1)}. \quad (37)$$

To evaluate the spectrum of the theory we need to compute the period integrals (8), with a similar expression for $\tilde{a}_j$ with the corresponding contour γ_j . The Seiberg-Witten




Figure 2: Diagram of the branch points and cycles for the single hypermultiplet case.

form $d\lambda$ for the curve (32) is given by

$$d\lambda = -\frac{X^{N+1}dX}{i2\pi Y} \left(\frac{1}{2(X+m)} - \frac{N}{X} \right). \quad (38)$$

Inserting the parametrization (36) we find the following expansion in powers of (m/Λ) for the periods:

$$\begin{aligned} \tilde{a}_j &= \frac{\Lambda}{4\pi} \left(e^{i4\pi j/(2N-1)} G(N+1) + \frac{m}{(2N-1)\Lambda} G(N) + \mathcal{O}((m/\Lambda)^2) \right) \\ a_{D_k} &= \tilde{a}_j|_{j \rightarrow k+1/2}, \end{aligned} \quad (39)$$

valid for $1 \leq j \leq (N-1)$ and $1 \leq k \leq (N-2)$. The coefficients $G(n)$ are given by

$$\begin{aligned} G(n) &:= 4 \int_0^\pi d\varphi \frac{e^{i\varphi(2n-1)/(2N-1)}}{\sqrt{e^{i2\varphi} - 1}} \\ &= -i4\pi^{3/2} \frac{e^{i\pi(n-1/2)/(2N-1)}}{\Gamma(\frac{1}{2} + \frac{n-1/2}{2N-1}) \Gamma(1 - \frac{n-1/2}{2N-1})}. \end{aligned} \quad (40)$$

The remaining period $a_{D_{N-1}}$ is given by the integral

$$a_{D_{N-1}} = \frac{1}{i\pi} \int_{X_{2N}}^{X_{2N-1}} \frac{X^{N+1}dX}{Y} \left(\frac{1}{2(X+m)} - \frac{N}{X} \right). \quad (41)$$

The leading order contribution is straightforward to obtain, one simply sets $m = 0$ and reduces the integral to

$$a_{D_{N-1}} = i \frac{N-1/2}{\pi} \int_0^\Lambda \frac{X^{N-1/2}dX}{\sqrt{X^{2N}-1}}$$

$$\begin{aligned}
&= \frac{\Lambda}{\sqrt{2\pi}} \int_0^\infty dx \frac{e^{-(1/2+1/(N-1/2))x}}{\sqrt{\sinh(x)}} \\
&= \frac{\Lambda}{2\sqrt{\pi}} \frac{\Gamma(\frac{1}{2} + \frac{1}{2N-1})}{\Gamma(1 + \frac{1}{2N-1})}
\end{aligned} \tag{42}$$

where in the second line we have used the change of variables $(X/\Lambda)^{N-1/2} := e^x$. A consistency check on the periods arises by noting that the a_{D_k} 's must satisfy

$$\sum_{k=0}^{N-1} a_{D_k} = m. \tag{43}$$

This constraint arises by noting that the sum of contours $\sum_{k=0}^{N-1} \alpha_k$ can be deformed to a contour encircling the pole of $d\lambda$ at $X = -m$ and the pole at infinity. The residue of $d\lambda$ at both poles is simply $m/2$ leading to the right-hand-side of (43). One can indeed show that the leading order piece to the periods of $\mathcal{O}(\Lambda)$ sums to zero, as it must. At $\mathcal{O}(m)$ one finds that

$$\sum_{k=0}^{N-2} a_{D_k} = \frac{N-1}{2N-1} m. \tag{44}$$

This implies that the $\mathcal{O}(m)$ contribution to the $a_{D_{N-1}}$ period must be $m(1/2 + 1/(2(2N-1)))$. A more careful evaluation of this period indeed shows that this is the case. Moreover we have checked that the sum in (43) cancels at $\mathcal{O}(m^2)$ and $\mathcal{O}(m^3)$ to $\mathcal{O}(1/N)$ and $\mathcal{O}(1)$ respectively. The period $a_{D_{N-1}}$ is given to $\mathcal{O}(m^3)$ in appendix B.

The derivatives of the periods obtained in a similar manner, and are given by

$$\begin{aligned}
\frac{\partial \tilde{a}_j}{\partial \xi_n} &= -\frac{\Lambda^{n-N}}{2\pi(2N-1)} e^{i2\pi j(n-1/2)/(N-1/2)} \left(G(n) - \frac{m}{\Lambda} \left(1 - \frac{n-1}{2N}\right) e^{-i2\pi j/(N-1/2)} G(n-2N) \right. \\
&\quad \left. + \dots \right), \quad 1 \leq j \leq (N-1) \\
\frac{\partial a_{D_k}}{\partial \xi_n} &= \frac{\partial \tilde{a}_j}{\partial \xi_n} \Big|_{j \rightarrow (k+1/2)}, \quad 1 \leq k \leq (N-2)
\end{aligned} \tag{45}$$

where both expressions are valid for $1 \leq n \leq (N-1)$. The remaining periods are given to leading order by

$$\frac{\partial a_{D_{N-1}}}{\partial \xi_n} = -\frac{\Lambda^{n-N}}{n\sqrt{\pi}} \frac{\Gamma(1 + \frac{n}{2N-1})}{\Gamma(\frac{1}{2} + \frac{n}{2N-1})}. \tag{46}$$

We have not been able to find an explicit analytic form for the period matrix given the formulae above. While in principle constructing the leading order piece of the period matrix should proceed as in the two-hypermultiplet case, we have not been able to invert the matrix $\partial \tilde{a}_k / \partial \xi_n$ analytically.

5 Probe metric

To compute the probe metric we should in principle consider the full $SU(N) \times U(n_3)$ gauge theory dynamics. Since we wish to treat the D3-branes as probe branes, we will take a limit

where the dynamics of the $U(n_3)$ is frozen out, and the theory reduces to $SU(N)$ coupled to n_3 hypermultiplets. Nevertheless, we obtain the probe metric from the prepotential of this theory by treating the hypermultiplet masses m_f as vacuum expectation values of scalars in vector multiplets on the same footing as the a_i . The probe metric is then

$$ds_{probe}^2 = \frac{d^2 \mathcal{F}}{dm^2} dm^2 , \quad (47)$$

where the derivative of $\mathcal{F}$ is defined as

$$d\mathcal{F} = \frac{\partial \mathcal{F}}{\partial m} dm + \sum_{i=1}^N \frac{\partial \mathcal{F}}{\partial a_i} da_i . \quad (48)$$

Fortunately, D'Hoker et al. [12] have obtained a closed form expression for the prepotential

$$2\mathcal{F} = \sum_{i=1}^{N-1} a_i a_{Di} + \sum_{f=1}^{n_3} m_{Df} m_f + \frac{1}{2\pi i} (Res_{P_+}(zd\lambda) Res_{P_+}(z^{-1}d\lambda) + Res_{P_-}(zd\lambda) Res_{P_-}(z^{-1}d\lambda)), \quad (49)$$

where

$$m_{Df} = \int_{P_f}^{P_-} d\lambda , \quad (50)$$

and P_f are the positions of the poles in $d\lambda$, and $P_{\pm}$ is infinity on the sheet where $y = \pm\sqrt{A^2 - B}$. The notation $Res_P(zd\lambda)$ denotes the residue of the form $zd\lambda$ at the pole P . The form $d\lambda$ and the function z will be defined below.

To make these m_D integrals well defined, a regularization procedure is needed. To set this up in a coordinate invariant way, the abelian integral $E = \log(Y + A)$ is introduced, with¹ $d\lambda = XdE$. The integrals of $d\lambda$ are then uniquely defined by imposing the following asymptotic conditions: near the P_f , $z = e^{-E}$,

$$\lambda(z) = -m_f \log z + \lambda(P_f) + \mathcal{O}(z) , \quad (51)$$

near P_- , $z = e^{\frac{E}{N-n_3}}$,

$$\lambda(z) = -Res_{P_-}(zd\lambda) + \mathcal{O}(z) \quad (52)$$

and near P_+ , $z = e^{-\frac{E}{N}}$,

$$\lambda(z) = -Res_{P_+}(zd\lambda) + \mathcal{O}(z) . \quad (53)$$

Computing the prepotential using (49) is now straightforward in principle given the results of the previous sections. In practice however evaluating the necessary integrals to the relevant order is tedious. The period integrals along the enhancon circle were given in the previous sections to linear order in the hypermultiplet masses. To obtain higher

¹This $d\lambda$ agrees with our previous expression in (9) up to the additive piece $XdXB'/(2B)$ which does not contribute to the a_j and a_{Dk} integrals. Rather this term only contributes to the m_D integrals, hence we do not change notation here.

order corrections one need only expand the integrand to higher order in the masses. The integrals can then be done exactly as discussed for the lower order contributions. The primary source of difficulty is the $a_{D_{N-1}}$ and m_D integrals (in either the single or double hypermultiplet case). We sketch the computation of these integrals in the appendix and give their results to $\mathcal{O}(m^3)$. Before discussing the limit of $m \ll \Lambda$ relevant for probing the enhancon, we note that in the perturbative limit $m \gg \Lambda$, we recover the supergravity result (3) provided we make the identifications $\rho_7 = \Lambda$. If we also wish the probe D7-brane computation [2] match with the supergravity result, we must identify $\rho_3 = \Lambda$. Subleading $1/N$ corrections also appear, with a form qualitatively the same as the D7-brane probe case described in [2].

In the limit $m \ll \Lambda$, we find

$$ds_{probe}^2 = \mathcal{O}(Nm^0) + \mathcal{O}(N^0 \frac{m}{\Lambda}) + \mathcal{O}(N^0 (\frac{m}{\Lambda})^2) + \dots \quad (54)$$

and all terms appear as an expansion in m/Λ with order one coefficients. The $\mathcal{O}(Nm^0)$ comes purely from the second residue term appearing in the formula for the prepotential (49). This term however is necessary in order to recover the correct perturbative form of the prepotential.

Computing the curvature of the probe metric in the form of the Ricci scalar, we find

$$R \sim \frac{\Lambda^2}{N(\log(|m|/\Lambda))^3 |m|^2} \quad (55)$$

when $|m| \gg \Lambda$. At a typical point, the curvature is of order $1/N$, and becomes of order one near the enhancon shell.

When $m \ll \Lambda$

$$R \sim \mathcal{O}(N^{-2}m^0) + \dots \quad (56)$$

This is the one of the main results of this paper. This result is consistent with the proposed flat interior for the enhancon shell of [1] as it does reduce to flat geometry as $N \rightarrow \infty$. In obtaining this result it was crucial to include the residue terms in the formula for the prepotential (49). Without these terms the $\mathcal{O}(m^0)$ piece of the metric (54) would have come with an $\mathcal{O}(1)$ coefficient, so that the curvature would also have been $\mathcal{O}(1)$.

In [2] we considered a D7 probe near the enhancon shell, where critical behavior emerges. This limit has also been studied by Ferrari [3, 4]. It would be interesting to study the analogous limit for the D3 probe, but we leave that for the future.

6 Conclusions

In this paper we have explored the enhancon mechanism from the point of view of $\mathcal{N} = 2$ supersymmetric $SU(N)$ gauge theory with fundamental hypermultiplets. We have obtained the exact low energy effective action for this theory, including non-perturbative effects. Our earlier results [2] explored the moduli space metric of the pure $SU(N)$ gauge theory, including configurations with probe branes both far and near the enhancon shell.

In the present work, we studied the enhancon in the presence of D3 brane probes which allowed us to probe the region inside the enhancon shell. In particular, we were able to show that at large N the geometry inside the enhancon shell is flat, i.e., curvature appears at order $1/N^2$. Our results support the supergravity picture of the enhancon resolution mechanism advocated in [1], and improve upon it in the gauge theory limit by including the full dynamics of the brane sources.

The gauge theory description makes clear the special role played by spherical symmetry in this resolution of the repulson singularity. The enhancon is at best marginally stable, since there exist exactly flat directions in moduli space corresponding to the quantum corrected position coordinates a_i of the D7-branes. From the gauge theory perspective, one can easily construct configurations where multiple branch cuts collide in the curve (4), though these points in moduli space will generally be non-spherically symmetric. These degenerations give rise to extra light particles creating singularities in the moduli space metric. Some of these degenerations are related to conformal field theories. From the supergravity viewpoint, these should be dual to nonsingular throats opening up in the geometry of the form anti-de Sitter space cross a sphere. Others are related to Argyres-Douglas points, and it remains unknown how these singularities will be resolved from the supergravity viewpoint. One possibility is there is no satisfactory resolution of these singularities purely within supergravity, and one must look for a more intrinsically stringy method of singularity resolution [13].

Acknowledgements

We thank G.L. Alberghi for collaboration during the early stages of this work. We thank E. D'Hoker for a helpful comment. This research is supported in part by DOE grant DE-FE0291ER40688-Task A.

A Appendix

In this appendix we collect some facts concerning the series expansions of various quantities in powers of the dimensionless quantities m_1/Λ and m_2/Λ for the two hypermultiplet case studied in (3). To begin with we consider the roots of the Seiberg-Witten curve,

$$\tilde{Y}^2 = \tilde{X}^{2N} - (\tilde{X} + \tilde{m})(\tilde{X} + \xi\tilde{m}), \quad (57)$$

rewritten in terms of dimensionless variables $\tilde{Y} \equiv Y/\Lambda^{2N}$, $\tilde{X} \equiv X/\Lambda$, $\tilde{m} \equiv m/\Lambda$, and $\xi = m_2/m_1$ with $m = m_1$. We search for roots of the polynomial by demanding that solutions of the form

$$\tilde{X} = \tilde{X}^{(0)}(1 + \sum_{j=1}^{\infty} \tilde{m}^j \tilde{X}^{(j)}) \quad (58)$$

hold order by order in the expansion parameter $\tilde{m}$. To zeroth order this will yield the solutions found previously in (6) for $X_k^{(0)}$. The coefficient of the $\tilde{m}^j$ term in the polynomial will then yield a linear equation for $\tilde{X}^{(j)}$ in terms of the $\tilde{X}^{(l)}$'s for $0 \leq l \leq (j-1)$. Consider in particular the case where $\tilde{X}^{(0)} = e^{i\pi k/(N-1)}$. It follows by induction that

$\tilde{X}^{(j)} \propto e^{-i\pi jk/(N-1)}$ where the coefficient of proportionality is independent of k . Denoting this coefficient by $d_j(\xi)$ we find the expansion for the roots

$$\tilde{X}_k = e^{i\pi k/(N-1)} \sum_{j=0}^{\infty} d_j(\xi) \tilde{m}^j e^{-i\pi jk/(N-1)}. \quad (59)$$

Now we can use this expansion to prove the statement in (3) that the series expansion of $\tilde{Y}$ (working with dimensionless quantities now) is given simply by (11). Recall that we substitute for $\tilde{X}$ in $\tilde{Y}$ the expansion $\tilde{X}_k|_{k \rightarrow k+\varphi/\pi}$. Expanding $\tilde{Y}$ in powers of $\tilde{m}$, one obtains exactly the same coefficients for each power $\tilde{m}^j$ as above when solving for the roots, providing that these coefficients are expressed in terms of the $\tilde{X}^{(j)}$'s and not their explicit expressions. The only exception to this occurs for $j = 0$. It follows immediately that the coefficients of the $\tilde{m}^j$ terms vanish by construction for $j \geq 1$. On the other hand the $j = 0$ term yields precisely (11).

A simple consequence of the expansion of the roots (59) and the parametrization (10) is the expansion of the periods given in (14). This follows easily by inserting the expansion (59) evaluated at $k \rightarrow k + \varphi/\pi$ into the Seiberg-Witten period $d\lambda$ and using the expansion of Y just derived.

B Appendix

In this appendix we present the results of the $a_{D_{N-1}}$ and m_D^{bare} period computations in both the single and double hypermultiplet cases and sketch briefly how these expressions were derived. For the single hypermultiplet case we find (setting $\Lambda = 1$ here and in the remaining formulae in this appendix)

$$\begin{aligned} a_{D_{N-1}} &= \frac{1}{2\sqrt{\pi}} \frac{\Gamma(\frac{1}{2} + \frac{1}{2N-1})}{\Gamma(1 + \frac{1}{2N-1})} + \frac{m}{2} \left(1 + \frac{1}{2N-1}\right) \\ &+ \frac{m^2}{4\sqrt{\pi}(2N-1)} \frac{\Gamma(\frac{1}{2} - \frac{1}{2N-1})}{\Gamma(1 - \frac{1}{2N-1})} \left(1 + \frac{1}{2N-1}\right) \\ &- \frac{m^3}{6\sqrt{\pi}(2N-1)} \frac{\Gamma(\frac{1}{2} - \frac{2}{2N-1})}{\Gamma(1 - \frac{2}{2N-1})} \left(1 - \frac{9}{2N-1} + \frac{2}{(2N-1)^2}\right) \\ &\approx \frac{1}{2} + \frac{m}{2}((-1)^N + \frac{1}{2N-1}) + \frac{m^2}{8N} - \frac{m^3}{12N} \end{aligned} \quad (60)$$

and

$$\begin{aligned} m_D^{bare} &= \frac{N-1}{i2\pi} X_{\infty} + \frac{m}{i2\pi} \ln(X_{\infty}) - \frac{m \ln(\delta m/m^N)}{i2\pi} + m \left(\frac{\ln(2) + 1}{i2\pi} - \frac{(i2 \ln(2) - \pi)}{4\pi(2N-1)} \right) \\ &- \frac{(-1)^{1/2-1/(2N-1)}}{8\pi^{3/2}} m^2 \Gamma\left(\frac{1}{2} - \frac{1}{2N-1}\right) \left(\Gamma\left(2 + \frac{1}{2N-1}\right) + \frac{2}{2N-1} \Gamma\left(1 + \frac{1}{2N-1}\right) \right. \\ &\left. + \Gamma\left(\frac{1}{2N-1}\right) \right) + \frac{(-1)^{1/2-2/(2N-1)}}{4\pi^{3/2}} m^3 \Gamma\left(\frac{1}{2} - \frac{2}{2N-1}\right) \left(\frac{1}{6} \Gamma\left(3 + \frac{2}{2N-1}\right) \right. \end{aligned} \quad (61)$$

$$\begin{aligned}
& + \frac{1}{2N-1} \frac{1}{2} \left(\Gamma\left(2 + \frac{2}{2N-1}\right) + \Gamma\left(1 + \frac{2}{2N-1}\right) + \Gamma\left(\frac{2}{2N-1}\right) \right) \\
& \approx \frac{N-1}{i2\pi} X_\infty + \frac{m}{i2\pi} \ln(X_\infty) - \frac{m \ln(\delta m/m^N)}{i2\pi} + m \frac{\ln(2)+1}{i2\pi} - i \frac{3}{8\pi} m^2 + i \frac{5}{24\pi} m^3,
\end{aligned}$$

where we have given both the exact expression to cubic order in m and the large N approximate form. Similarly for the double hypermultiplet case we find

$$\begin{aligned}
a_{D_{N-1}} &= \frac{N-1}{N\sqrt{\pi}} \frac{\Gamma(\frac{3}{2} + \frac{1}{2(N-1)})}{\Gamma(1 + \frac{1}{2(N-1)})} + \frac{m^2}{\sqrt{\pi}(N-2)} \Gamma\left(\frac{3}{2} - \frac{1}{2(N-1)}\right) \left(\frac{2(N-1)}{\Gamma(-1 - \frac{1}{2(N-1)})} \right. \\
&+ \left. \frac{N+1}{\Gamma(-\frac{1}{2(N-1)})} - \frac{1}{\Gamma(1 - \frac{1}{2(N-1)})} \right) \\
&+ \frac{m^3}{\sqrt{\pi}(N-3)} \Gamma\left(\frac{3}{2} - \frac{1}{N-1}\right) \left(\frac{8(N-1)}{3\Gamma(-2 - \frac{1}{N-1})} + \frac{2(2N+1)}{\Gamma(-1 - \frac{1}{N-1})} \right. \\
&- \left. \frac{1}{\Gamma(-\frac{1}{N-1})} + \frac{1}{\Gamma(1 - \frac{1}{N-1})} \right) \\
&\approx \frac{1}{2} - \frac{m^2}{4N} - \frac{m^3}{6N}
\end{aligned} \tag{62}$$

and

$$\begin{aligned}
m_D^{\text{bare}} &= \frac{1}{i2\pi} \left(-(N-2)X_\infty - 2m \ln(X_\infty) + 2m \ln(\delta m/m^{N/2}) \right) \\
&- \frac{(-1)^{1/(2(N-1))}}{\sqrt{\pi}} \frac{N-1}{N} \Gamma\left(\frac{3}{2} + \frac{1}{2(N-1)}\right) \Gamma\left(-\frac{1}{2(N-1)}\right) + m \left(2 - 2 \ln(2) + \frac{i\pi}{2} \right) \\
&- m^2 \frac{(-1)^{1/(2(N-1))}}{2\sqrt{\pi}} \frac{N}{(N-1)(N-2)} \Gamma\left(\frac{3}{2} - \frac{1}{2(N-1)}\right) \Gamma\left(\frac{1}{2(N-1)}\right) \\
&+ m^3 \frac{(-1)^{-1/(N-1)}}{3\sqrt{\pi}} \frac{N(7N-11)}{(N-1)^2(N-3)} \Gamma\left(\frac{3}{2} - \frac{1}{N-1}\right) \Gamma\left(\frac{1}{N-1}\right) \\
&= \frac{1}{i2\pi} \left(-(N-2)X_\infty - 2m \ln(X_\infty) + 2m \ln(\delta m/m^{N/2}) + N \right. \\
&+ \left. m \left(2 - 2 \ln(2) + \frac{i\pi}{2} \right) - \frac{m^2}{2} + \frac{7m^3}{6} \right).
\end{aligned} \tag{63}$$

For concreteness we shall now briefly sketch the derivation of $a_{D_{N-1}}$ for the single hypermultiplet. The remaining periods follow via identical reasoning however. The integral that we need is given by

$$a_{D_{N-1}} = \oint_{\alpha_{N-1}} d\lambda = \int_{X_0}^{X_{2N}} d\lambda \tag{64}$$

where $d\lambda$ is given by (9). To evaluate the integral we shall split the integration region into four parts: (i) $(X_{2N} + m) \leq (X + m) \leq \xi_0 m^{2N}$, (ii) $\xi_0 m^{2N} \leq (X + m) \leq \xi_1 m$, (iii) $\xi_1 m \leq (X + m) \leq (X_0 + m - \xi_2 m/N)$, and (iv) $(X_0 - \xi_2 m/N) \leq X \leq X_0$. The parameters ξ_j are all numbers of $\mathcal{O}(10)$. In region (i) we make the change of integration

variable to $X = -m + um^{2N}$. The integrand can then be series expanded in m and the resulting integrals are straightforward to evaluate (at least to the order presented above). In region (ii) one series expands the integrand in the variable $(X + m)$. In region (iii) one series expands the integrand in m , and in region (iv) one defines the variable of integration $X = X_0 - vm/N$ and then series expands in m . It is straightforward to check that these series expansions are valid over the ranges given above. As a consistency check one notes that the full integral must be independent of the (relatively) arbitrary parameters ξ_j introduced in the intermediate steps. Indeed one can show that this cancellation takes place to the order that we are working.

References

- [1] Clifford V. Johnson, Amanda W. Peet, and Joseph Polchinski. Gauge theory and the excision of repulson singularities. *Phys. Rev.*, D61:086001, 2000, hep-th/9911161.
- [2] Gian Luigi Alberghi, Steven Corley, and David A. Lowe. Moduli space metric of $n=2$ supersymmetric $su(n)$ gauge theory and the enhancon. 2002, hep-th/0204050.
- [3] Frank Ferrari. The large n limit of $n = 2$ super yang-mills, fractional instantons and infrared divergences. *Nucl. Phys.*, B612:151–170, 2001, hep-th/0106192.
- [4] Frank Ferrari. Exact amplitudes in four dimensional non-critical string theories. *Nucl. Phys.*, B617:348–364, 2001, hep-th/0107096.
- [5] Clifford V. Johnson, Robert C. Myers, Amanda W. Peet, and Simon F. Ross. The enhancon and the consistency of excision. *Phys. Rev.*, D64:106001, 2001, hep-th/0105077.
- [6] N. Seiberg and Edward Witten. Electric - magnetic duality, monopole condensation, and confinement in $n=2$ supersymmetric yang-mills theory. *Nucl. Phys.*, B426:19–52, 1994, hep-th/9407087.
- [7] N. Seiberg and Edward Witten. Monopoles, duality and chiral symmetry breaking in $n=2$ supersymmetric qcd. *Nucl. Phys.*, B431:484–550, 1994, hep-th/9408099.
- [8] Philip C. Argyres and Michael R. Douglas. New phenomena in $su(3)$ supersymmetric gauge theory. *Nucl. Phys.*, B448:93–126, 1995, hep-th/9505062.
- [9] Amihay Hanany and Yaron Oz. On the quantum moduli space of vacua of $n=2$ supersymmetric $su(n(c))$ gauge theories. *Nucl. Phys.*, B452:283–312, 1995, hep-th/9505075.
- [10] Philip C. Argyres and Alon E. Faraggi. The vacuum structure and spectrum of $n=2$ supersymmetric $su(n)$ gauge theory. *Phys. Rev. Lett.*, 74:3931–3934, 1995, hep-th/9411057.

- [11] A. Klemm, W. Lerche, S. Yankielowicz, and S. Theisen. Simple singularities and $n=2$ supersymmetric yang-mills theory. *Phys. Lett.*, B344:169–175, 1995, hep-th/9411048.
- [12] Eric D'Hoker, I. M. Krichever, and D. H. Phong. The renormalization group equation in $n = 2$ supersymmetric gauge theories. *Nucl. Phys.*, B494:89–104, 1997, hep-th/9610156.
- [13] David A. Lowe. Renormalization group flows from gravity in anti-de sitter space versus black hole no-hair theorems. *JHEP*, 04:011, 2000, hep-th/0003042.